

# BIRTH DEFECT RISK FACTOR SERIES:

## DIAPHRAGMATIC HERNIA

### DEFINITION

---

Congenital diaphragmatic hernia occurs when the diaphragm is incompletely formed and the abdominal contents herniate into the chest. Infants with diaphragmatic hernia tend to have other additional birth defects, including chromosomal abnormalities (Forrester and Merz, 1998; Robert et al., 1997; Cannon et al., 1996; Howe et al., 1996; Langham et al., 1996; Torfs et al., 1992; Wenstrom et al., 1991; Adzick et al., 1985; David and Illingworth, 1976). Most diaphragmatic hernias occur on the left side of the body (Forrester and Merz, 1998; Robert et al., 1997; Howe et al., 1996; Torfs et al., 1992; Rasheed et al., 1992; Wenstrom et al., 1991; Adzick et al., 1985; David and Illingworth, 1976).

Diaphragmatic hernia can be detected prenatally with ultrasonography (Langham et al., 1996; Vintzileos et al., 1987). As a result, prenatal diagnosis and elective termination can reduce the birth prevalence of diaphragmatic hernia (Forrester and Merz, 1998; Riley et al., 1998; Cannon et al., 1996; Howe et al., 1996; Stoll et al., 1995; Juilian-Reynier et al., 1994; Stoll et al., 1992; Torfs et al., 1992; Adzick et al., 1985).

### EMBRYOLOGY

---

The diaphragm develops from four different structures: the septum transversum, pleuroperitoneal membranes, dorsal mesentery of the esophagus, and the body wall. The septum transversum is embryonic mesoderm that separates the pericardial (heart) cavity from the gastrointestinal system. The pleuroperitoneal membranes separate the pleural (lung) and the peritoneal (abdominal) cavities. The dorsal mesentery of the esophagus is a layer of peritoneum. The septum transversum, pleuroperitoneal membranes, and dorsal mesentery of the esophagus fuse together by the sixth or seventh week of gestation to form the diaphragm. If the pleuroperitoneal membranes fail to fuse with the septum transversum and the dorsal mesentery of the esophagus by the tenth week of gestation, diaphragmatic hernia results. Diaphragmatic hernia is more common on the left side because the right pleuroperitoneal membrane usually fuses before the left pleuroperitoneal membrane because of the embryonic liver on the right side of the body.

### DEMOGRAPHIC AND REPRODUCTIVE FACTORS

---

Several studies have investigated the relationship of **race/ethnicity** and risk for diaphragmatic hernia and failed to find any significant effect (Forrester and Merz, 1998; Robert et al., 1997; Langham, 1996; Torfs et al., 1992). However, one study did report diaphragmatic hernia to be more common among whites than nonwhites (Yang et al., 1994), and another reported the rate of diaphragmatic hernia to be higher in Europeans and South Asians than in Caribbeans (Leck and Lancashire, 1995).

**Secular trends** have been reported by several investigations (Forrester and Merz, 1998; Yang et al., 1994; Torfs et al., 1992). However, the trends were neither consistent nor statistically significant. One study observed **seasonal variation** in diaphragmatic hernia rates; however, this variation differed between types of diaphragmatic hernia (Torfs et al., 1992). Moreover, other investigations failed to identify any seasonal variation (Castilla et al., 1990; Bound et al., 1989).

Prevalence of diaphragmatic hernia shows variation by **geographic location**, with the rate being higher in rural than urban areas (Forrester and Merz, 1998; Torfs et al., 1992). One investigation failed to identify any association between diaphragmatic hernia and **altitude** (Castilla et al., 1999).

**Maternal age** does not appear to influence risk for diaphragmatic hernia (Forrester and Merz, 1998; Robert

et al., 1997; Torfs et al., 1992; David and Illingworth, 1976), although one study did report an increased risk with increased maternal age (Hollier et al., 2000). No clear association between diaphragmatic anomalies and **paternal age** has been identified (McIntosh et al., 1995). **Parity** does not affect diaphragmatic hernia rates (Robert et al., 1997; Torfs et al., 1992; David and Illingworth, 1976). The **recurrence risk** of a woman having another infant with diaphragmatic hernia has been reported to be 0.9-2% (Buyse, 1990).

**Infant sex** is associated with diaphragmatic hernia risk. Males were more likely than females to have the defect (Lary and Paulozzi, 2001; Robert et al., 1997; Yang et al., 1994; Torfs et al., 1992; Rasheed et al., 1992; Buyse, 1990; Rose-Spencer et al., 1981). Diaphragmatic hernia has been associated with lower **birth weight** and **intrauterine growth retardation** (Riley et al., 1998; Mili et al., 1991; Khoury et al., 1988). No association was found between anomalies of the diaphragm and **macrosomia** (Lapunzina et al., 2002; Waller et al., 2001). The literature on the relationship between diaphragmatic hernia and **multiple gestation pregnancy** has been inconsistent (Mastroiacovo, 1999; Riley et al., 1998; Robert et al., 1997; Torfs et al., 1992; Doyle et al., 1991; Kallen, 1986).

One investigation found no association between parental **consanguinity** and diaphragmatic hernia (Rittler et al., 2001).

## FACTORS IN LIFESTYLE OR ENVIRONMENT

---

Several investigations examined various maternal and environmental factors in relation to diaphragmatic hernia and found no increased risk for **maternal thyroid dysfunction (hyperthyroidism, hypothyroidism)** or **occupational and environmental chemicals** (Bos et al., 1994; Khoury et al., 1989). Studies have reported no association between **cephalosporin antibiotics, augmentin, ampicillin, calcium channel blockers**, or the **benzodiazepines** nitrazepam, medazepam, tofisopam, alprazolam, and clonazepam and anomalies of the diaphragm (Eros et al., 2002; Czeizel et al., 2001a; Czeizel et al., 2001b; Czeizel et al., 2001c; Sorensen et al., 2001). A **herbicide Nitrofen** has been found to induce diaphragmatic hernia in rats (Bos et al., 1994). Another study reported a higher than expected diaphragmatic hernia prevalence among infants whose mothers had **epilepsy**, although this increased risk may be associated with the **anticonvulsants** the mothers may have been taking (Bertollini et al., 1985).

An investigation failed to identify any significant association between diaphragmatic hernia and proximity to various types of **industry** (Castilla et al., 2000). One study observed no association between diaphragmatic hernia and **water chlorination** (Kallen and Robert, 2000).

No association between maternal **folic acid** use and diaphragmatic defects has been reported (Czeizel et al., 1996). Furthermore, a study that examined **co-trimoxazole**, a combination of trimethoprim and sulfamethoxazole that is a folic acid antagonist, failed to find any association between the medication and diaphragmatic anomalies (Czeizel, 1990).

## REFERENCES

---

Adzick NS, Harrison MR, Glick PL, Nakayama DK, Manning FA, deLorimier AA. Diaphragmatic hernia in the fetus: Prenatal diagnosis and outcome in 94 cases. *J Pediatr Surg* 1985;20:357-361.

Bertollini R, Mastroiacovo P, Segni G. Maternal epilepsy and birth defects: a case-control study in the Italian Multicentric Registry of Birth Defects (IPIMC). *Eur J Epidemiol* 1985;1:67-72

Bos AP, Pattenier AM, Grobbee RE, Lindhout D, Tibboel D, Molenaar JC. Etiological aspects of congenital diaphragmatic hernia: results of a case comparison study. *Hum Genet* 1994;94:445-446.

Bound JP, Harvey PW, Francis BJ. Seasonal prevalence of major congenital malformations in the Fylde of Lancashire 1957-1981. *J Epidemiol Community Health* 1989;43:330-342.

Buyse ML, editor-in-chief. *Birth Defect Encyclopedia*. Cambridge, Massachusetts: Blackwell Scientific Publications, 1990.

Cannon C, Dildy GA, Ward R, Varner MW, Dudley DJ. A population-based study of congenital diaphragmatic hernia in Utah: 1988-1994. *Obstet Gynecol* 1996;87:959-963.

Castilla EE, Campana H, Camelo JS. Economic activity and congenital anomalies: an ecologic study in Argentina. *Environ Health Perspect* 2000;108:193-197.

Castilla EE, Lopez-Camelo JS, Campana H. Altitude as a risk factor for congenital anomalies. *Am J Med Genet* 1999;86:9-14.

Castilla EE, Orioli IM, Lugarinho R, Dutra GP, Lopez-Camelo JS, Campana HE, Spagnolo A, Mastroiacovo P. Monthly and seasonal variations in the frequency of congenital anomalies. *Int J Epidemiol* 1990;19:399-404

Czeizel A. A case-control analysis of the teratogenic effects of co-trimoxazole. *Reprod Toxicol* 1990;4:305-313.

Czeizel AE, Toth M, Rockenbauer M. Population-based case control study of folic acid supplementation during pregnancy. *Teratology* 1996;53:345-351.

Czeizel AE, Rockenbauer M, Sorensen HT, Olsen J. Use of cephalosporins during pregnancy and in the presence of congenital abnormalities: a population-based, case-control study. *Am J Obstet Gynecol* 2001a;184:1289-1296.

Czeizel AE, Rockenbauer M, Sorensen HT, Olsen J. Augmentin treatment during pregnancy and the prevalence of congenital abnormalities. A population-based case-control teratologic study. *Eur J Obstet Gynecol Reprod Biol* 2001b;97:188-192.

Czeizel AE, Rockenbauer M, Sorensen HT, Olsen J. A population-based case-control teratologic study of ampicillin treatment during pregnancy. *Am J Obstet Gynecol* 2001c;185:140-147.

David TJ, Illingworth CA. Diaphragmatic hernia in the south-west of England. *J Med Genet* 1976;13:253-262.

Doyle PE, Beral V, Botting B, Wale CJ.

Congenital malformations in twins in England and Wales. *J Epidemiol Community Health* 1991;45:43-48.

Eros E, Czeizel AE, Rockenbauer M, Sorensen HT, Olsen J. A population-based case-control teratologic study of nitrazepam, medazepam, tofisopam, alprazolam and clonazepam treatment during pregnancy. *Eur J Obstet Gynecol Reprod Biol* 2002;101:147-154.

Forrester MB, Merz RD. Epidemiology of congenital diaphragmatic hernia, Hawaii, 1987-1996. *Hawaii Med J* 1998;57:586-589.

Hollier LM, Leveno KJ, Kelly MA, McIntire DD, Cunningham FG. Maternal age and malformations in singleton births. *Obstet Gynecol* 2000;96:701-706.

Howe DT, Kilby MD, Sirry H, Barker GM, Roberts E, Davison EV, McHugo J, Whittle MJ. Structural chromosome anomalies in congenital diaphragmatic hernia. *Prenat Diagn* 1996;16:1003-1009.

Julian-Reynier C, Philip N, Scheiner C, Auran Y, Chabal F, Maron A, Gombert A, Ayme S. Impact of prenatal diagnosis by ultrasound on the prevalence of congenital anomalies at birth in southern France. *J Epidemiol Community Health* 1994;48:290-296.

Kallen B. Congenital malformations in twins: a population study. *Acta Genet Med Gemellol (Roma)* 1986;35:167-178.

Kallen BA, Robert E. Drinking water chlorination and delivery outcome-a registry-based study in Sweden. *Reprod Toxicol* 2000;14:303-309.

Khoury MJ, Becerra JE, d'Almada PJ. Maternal thyroid disease and risk of birth defects in offspring: a population-based case-control study. *Paediatr Perinat Epidemiol* 1989;3:402-420.

Khoury MJ, Erickson JD, Cordero JF, McCarthy BJ. Congenital malformations and intrauterine growth retardation: a population study. *Pediatrics* 1988;82:83-90.

Langham MR, Kays DW, Ledbetter DJ, Frentzen B, Sanford LL, Richards DS. Congenital diaphragmatic hernia. Epidemiology and outcome. *Clin Perinatol* 1996;23:671-688.

Lapunzina P, Lopez Camelo JS, Rittler M, Castilla EE. Risks of congenital anomalies in

large for gestational age infants. J Pediatr 2002;140:200-204.

Lary JM, Paulozzi LJ. Sex differences in the prevalence of human birth defects: a population-based study. Teratology 2001;64:237-251.

Leck I, Lancashire RJ. Birth prevalence of malformations in members of different ethnic groups and in the offspring of matings between them, in Birmingham, England. J Epidemiol Community Health 1995;49:171-179.

Mastroiacovo P, Castilla EE, Arpino C, Botting B, Cocchi G, Goujard J, Marinacci C, Merlob P, Metneki J, Mutchinick O, Ritvanen A, Rosano A. Congenital malformations in twins: an international study. Am J Med Genet 1999;83:117-124.

McIntosh GC, Olshan AF, Baird PA. Paternal age and the risk of birth defects in offspring. Epidemiology 1995;6:282-288.

Mili F, Edmonds LD, Khoury MJ, McClearn AB. Prevalence of birth defects among low-birth-weight infants. A population study. Am J Dis Child 1991;145:1313-1318.

Rasheed K, Coughlan G, O'Donnell B. Congenital diaphragmatic hernia in the newborn. Outcome in 59 consecutive cases over a ten year period (1980-1989). Ir J Med Sci 1992;161:16-17

Riley MM, Halliday JL, Lumley JM. Congenital malformations in Victoria, Australia, 1983-95: an overview of infant characteristics. J Paediatr Child Health 1998;34:233-240.

Rittler M, Liascovich R, Lopez-Camelo J, Castilla EE. Parental consanguinity in specific types of congenital anomalies. Am J Med Genet 2001;102:36-43.

Robert E, Kallen B, Harris J. The epidemiology of diaphragmatic hernia. Eur J Epidemiol 1997;13:665-673

Rose-Spencer JA, Bloss RS, Beardmore HE. Congenital posterolateral diaphragmatic hernia: a retrospective study. Can J Surg 1981;24:515-517

Sorensen HT, Czeizel AE, Rockenbauer M, Steffensen FH, Olsen J. The risk of limb deficiencies and other congenital abnormalities in children exposed in utero to calcium channel blockers. Acta Obstet Gynecol Scand 2001;80:397-401.

Stoll C, Dott B, Alembik Y, Roth MP. Evaluation of routine prenatal diagnosis by a registry of congenital anomalies. Prenat Diagn 1995;15:791-800.

Stoll C, Alembik Y, Dott B, Roth MP, Finck S. Evaluation of prenatal diagnosis by a registry of congenital anomalies. Prenat Diagn 1992;12:263-270.

Torfs CP, Curry CJR, Bateson TF, Honore LH. A population-based study of congenital diaphragmatic hernia. Teratology 1992;46:555-565.

Vintzileos AM, Campbell WA, Nochimson DJ, Weinbaum PJ. Antenatal evaluation and management of ultrasonically detected fetal anomalies. Obstet Gynecol 1987;69:640-660.

Waller DK, Keddie AM, Canfield MA, Scheuerle AE. Do infants with major congenital anomalies have an excess of macrosomia? Teratology 2001;64:311-317.

Wenstrom KD, Weiner CP, Hanson JW. A five-year statewide experience with congenital diaphragmatic hernia. Am J Obstet Gynecol 1991;165:838-842.

Yang P, Khoury MJ, Stewart WF, Beaty TH, Chee E, Beatty JC, Diamond EL, Gordis L. Comparative epidemiology of selected midline congenital abnormalities. Genet Epidemiol 1994;11:141-154.

**Please Note:** The primary purpose of this report is to provide background necessary for conducting cluster investigations. It summarizes literature about risk factors associated with this defect. The strengths and limitations of each reference were not critically examined prior to inclusion in this report. Consumers and professionals using this information are advised to consult the references given for more in-depth information.

This report is for information purposes only and is not intended to diagnose, cure, mitigate, treat, or prevent disease or other conditions and is not intended to provide a determination or assessment of the state of health. Individuals affected by this condition should consult their physician and when appropriate, seek genetic counseling.